

(51) International Patent Classification <sup>6</sup> :

C07K 5/06, A23L 1/236

A1

(11) International Publication Number:

WO 99/12956

(43) International Publication Date:

18 March 1999 (18.03.99)

(21) International Application Number: PCT/US98/18902

(22) International Filing Date: 11 September 1998 (11.09.98)

(30) Priority Data:

60/058,505



11 September 1997 (11.09.97) US

(71) Applicant: THE NUTRASWEET COMPANY [US/US]; Suite 900, 200 World Trade Center, Chicago, IL 60654 (US).

(72) Inventors: PRAKASH, Indra; 4731 Sunflower Lane, Hoffman Estates, IL 60195 (US). WACHHOLDER, Kurt, L.; 1175 Price Drive, Elgin, IL 60120 (US).

(74) Agents: MANDRA, Raymond, R. et al.; Fitzpatrick, Cella, Harper &amp; Scinto, 30 Rockefeller Plaza, New York, NY 10112-3801 (US).

(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

**Published***With international search report.**Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.*(54) Title: ACID SALTS OF N-[N-(3,3-DIMETHYLBUTYL)-L- $\alpha$ -ASPARTYL]-L-PHENYLALANINE 1-METHYL ESTER**(57) Abstract**

Dipeptide sweeteners are disclosed that are salts of N-[N-(3,3-dimethylbutyl)-L- $\alpha$ -aspartyl]-L-phenylalanine 1-methyl ester represented by formula (I) wherein X<sup>-</sup> is selected from the group consisting of Cl<sup>-</sup>, HSO<sub>4</sub><sup>-</sup>, H<sub>2</sub>PO<sub>3</sub><sup>-</sup>, citrate, HCO<sub>3</sub><sup>-</sup>, fumarate, malate, maleate, tartarate, acetate, benzoate or gluconate. Also disclosed is a liquid low-calorie sweetener container such salts.

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Larvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

- 1 -

# TITLE

ACID SALTS OF N-[N-(3,3-DIMETHYLBUTYL)-  
L- $\alpha$ -ASPARTYL]-L-PHENYLALANINE 1-METHYL ESTER

5

## BACKGROUND OF THE INVENTION

### Field of the Invention

10 This invention relates to novel sweeteners. In particular, the invention relates to acid salts of the N-alkylated aspartame derivative, N-[N-(3,3-dimethylbutyl)-L- $\alpha$ -aspartyl]-L-phenylalanine 1-methyl ester, i.e., neotame. The invention also relates to a  
15 liquid low calorie sweetener containing such salts.


### Related Background Art

It is known that various N-substituted derivatives of  
20 aspartame, such as disclosed in U.S. Patent No. 5,480,668, are useful as sweetening agents. In particular, the N-alkylated aspartame derivative, N-[N-(3,3-dimethylbutyl)-L- $\alpha$ -aspartyl]-L-phenylalanine 1-methyl ester, is known as an extremely potent  
25 sweetening agent since its sweetening potency, on a weight basis, has been reported to be at least 50 times


- 2 -

that of aspartame and about 10,000 times that of sucrose.

- Since sweetening agents are often employed in aqueous solutions and beverages, it is important that they have an acceptable dissolution rate and an effective level of solubility to be commercially practicable. U.S. Patent Nos. 4,029,701 and 4,031,258 describe inorganic salts of dipeptides that provide improved solubility in aqueous solution while maintaining the sweet property of the dipeptide. U.S. Patent No. 4,448,716 describes certain dipeptide sweetener-metal complexes that provide improved dissolution and solubility. U.S. Patent 4,153,737 describes concentrated solutions of dipeptide salts in a non-aqueous system (concentrated liquid low calorie sweetener). N-[N-(3,3-dimethylbutyl)-L- $\alpha$ -aspartyl]-L-phenylalanine 1-methyl ester, however, is not disclosed or suggested.
- Structurally, N-[N-(3,3-dimethylbutyl)-L- $\alpha$ -aspartyl]-L-phenylalanine 1-methyl ester and aspartame differ in that, in N-[N-(3,3-dimethylbutyl)-L- $\alpha$ -aspartyl]-L-phenylalanine 1-methyl ester, a bulky neohexyl substituent is present on the amine nitrogen.



Aspartame



Neotame

25

- This structural difference results in dramatic differences in the physical and chemical properties of these compounds. For example, the melting point of N-[N-(3,3-dimethylbutyl)-L- $\alpha$ -aspartyl]-L-phenylalanine 1-methyl ester is 80°C, while that of aspartame is 248°C.

30

- 3 -

In addition, N-[N-(3,3-dimethylbutyl)-L- $\alpha$ -aspartyl]-L-phenylalanine 1-methyl ester has much higher solubility in organic solvents than aspartame, and a much lower solubility in water. It is also known that N-[N-(3,3-dimethylbutyl)-L- $\alpha$ -aspartyl]-L-phenylalanine 1-methyl ester has a higher stability than aspartame under some pH conditions, as described in U.S. Patent No. 5,480,688. The pronounced difference in sweetness between the two compounds is further evidence of their chemical dissimilarity.

Moreover, it is also known that a primary amino group such as the one on aspartame (pKa 7.7) generally has a different pKa than those from a secondary amino group such as the one on N-[N-(3,3-dimethylbutyl)-L- $\alpha$ -aspartyl]-L-phenylalanine 1-methyl ester (pKa 8.1). Moreover, the pKa's of an amino acid are known to have a profound impact on food applications (Labuza, T.P. and Basisier, M.W., 1992, "Physical Chemistry of Foods", H.G. Schwartzber and R.W. Hartel (Eds.), Marcel Dekker, Inc., New York). It is also well known that a secondary amine group can not form Schiff base type compounds with carbonyl compounds while a primary amine may. Furthermore, N-[N-(3,3-dimethylbutyl)-L- $\alpha$ -aspartyl]-L-phenylalanine 1-methyl ester exhibits physiologically different behavior than aspartame as exemplified by the dramatic difference in sweetness. These differences are clearly indicative that the characteristics and properties of one can not be said to suggest those of the other.

While N-[N-(3,3-dimethylbutyl)-L- $\alpha$ -aspartyl]-L-phenylalanine 1-methyl ester is a highly potent sweetener, it is sparingly soluble in water and can give rise to dusting problems. Therefore, there is a need for N-[N-(3,3-dimethylbutyl)-L- $\alpha$ -aspartyl]-L-phenylalanine 1-methyl ester derivatives that have


- 4 -

good dissolution and solubility properties in aqueous systems, and avoid dusting problems often encountered with fine powders.

5

## SUMMARY OF THE INVENTION

This invention relates to dipeptide sweeteners that are salts of N-[N-(3,3-dimethylbutyl)-L- $\alpha$ -aspartyl]-L-phenylalanine 1-methyl ester possessing good  
 10 dissolution and solubility properties in aqueous systems. In particular, the salts of N-[N-(3,3-dimethylbutyl)-L- $\alpha$ -aspartyl]-L-phenylalanine 1-methyl ester of this invention are represented by the formula



15 wherein X<sup>-</sup> is selected from the group consisting of Cl<sup>-</sup>, HSO<sub>4</sub><sup>-</sup>, H<sub>2</sub>PO<sub>4</sub><sup>-</sup>, citrate, HCO<sub>3</sub><sup>-</sup>, fumarate, malate, maleate, tartarate, acetate, benzoate or gluconate. The invention is also related to a liquid low calorie sweetener containing the acid salts of this invention.

20

## BRIEF DESCRIPTION OF THE DRAWING

The figure is a graph comparing the aqueous dissolution of N-[N-(3,3-dimethylbutyl)-L- $\alpha$ -aspartyl]-L-phenylalanine 1-methyl ester at a target concentration  
 25 of 0.05% by weight with equivalent neotame concentrations, i.e., the concentration of the neotame delivered in each case is the same, of the phosphate and citrate salts of N-[N-(3,3-dimethylbutyl)-L- $\alpha$ -  
 30 aspartyl]-L-phenylalanine 1-methyl ester.

# DETAILED DESCRIPTION OF THE INVENTION

This invention is directed to salts of N-[N-(3,3-dimethylbutyl)-L- $\alpha$ -aspartyl]-L-phenylalanine 1-methyl ester, i.e., salts of neotame. U.S. Patent No. 5,480,668, U.S. Patent No. 5,510,508 and U.S. Patent No. 5,728,862, which describe the preparation of N-[N-(3,3-dimethylbutyl)-L- $\alpha$ -aspartyl]-L-phenylalanine 1-methyl ester are incorporated by reference herein as if fully set forth. Thus, the starting material may be readily prepared by one of ordinary skill in the art without undue experimentation.

The acid salts of this invention may be prepared by first slurrying the N-[N-(3,3-dimethylbutyl)-L- $\alpha$ -aspartyl]-L-phenylalanine 1-methyl ester in a solvent or mixture of solvents. Exemplary solvents may include water, methanol, acetone, tetrahydrofuran and the like. The slurry is stirred and an equimolar amount of the desired acid is slowly added to dissolve the N-[N-(3,3-dimethylbutyl)-L- $\alpha$ -aspartyl]-L-phenylalanine 1-methyl ester and form the salt of this invention. The salt may be recovered by evaporation of the solvent, freeze drying or spray drying the resulting solution. The order of the addition of sweetener to acid has not been found to be of consequence and can be readily determined by those skilled in the art. Acid salts of N-[N-(3,3-dimethylbutyl)-L- $\alpha$ -aspartyl]-L-phenylalanine 1-methyl ester prepared under these conditions do not show any racemization. Addition of excess of acid (more than 1 equivalent) does cause hydrolysis of the methyl ester.

The acids employed in the preparation of the salts of this invention are typically selected from compounds that have a pKa effectively lower than the pKa of the secondary amine of N-[N-(3,3-dimethylbutyl)-L- $\alpha$ -

- 6 -

aspartyl]-L-phenylalanine 1-methyl ester to result in the formation of the desired salt. Such compounds include, for example, hydrochloric acid, sulfuric acid, phosphoric acid, citric acid, fumaric acid, malic acid, 5 maleic acid, tartaric acid, acetic acid, benzoic acid, gluconic acid and carbonic acid. As such,  $X^-$  is a physiologically acceptable anion selected from the group consisting of  $Cl^-$ ,  $HSO_4^-$ ,  $H_2PO_4^-$ , citrate $^-$ ,  $HCO_3^-$ , fumarate, malate, maleate, tartarate, acetate, 10 benzoate or gluconate. These ions can be used alone or in combination.

Particularly preferred salts of this invention include the hydrochloride, phosphate, sulfate, citrate, 15 hydrobromide and carbonic salts of N-[N-(3,3-dimethylbutyl)-1-L- $\alpha$ -aspartyl]-L-phenylalanine 1-methyl ester.

It is believed that the acid salts of this invention 20 provide a number of improved properties over those of N-[N-(3,3-dimethylbutyl)-L- $\alpha$ -aspartyl]-L-phenylalanine 1-methyl ester. In particular, the aqueous solubility is increased and the dissolution rate of the composition is greatly improved. These acidic salts of 25 neotame are sweet and it is also believed that these salts have improved taste. Thus, these salts of N-[N-(3,3-dimethylbutyl)-L- $\alpha$ -aspartyl]-L-phenylalanine 1-methyl ester will be particularly useful in beverage systems, particularly since additional methods or 30 mechanical preparations are diminished or not necessary to provide for quick dissolution such as desired in a table top sweetener. The salts of this invention may be admixed with known bulking agents to prepare tablets, powdered or granular sweetener using methods 35 well known to those skilled in the art.



These salts may also be used to prepare a liquid, low-calorie sweetener by dissolving a high concentration of the salt of this invention in an aqueous or alcoholic system, e.g., water, ethanol, water/ethanol, propylene glycol or propylene glycol/water. Such a liquid, low-calorie sweetener may find utility in such foodstuffs as gelatin desserts, fruit flavored beverages, cereal, cake mixes, fruit juices, syrups, salad dressings, pet foods, carbonated soft drinks, table top sweeteners and the like. Such utilities are not restrictive since other applications may include cough medicines, tonics and the like. One embodiment of this invention of particular interest contemplates a liquid table top sweetener as a replacement for sucrose and other known sweeteners. The liquid low calorie sweetener generally will contain up to about 40% by weight of the salt of N-[N-(3,3-dimethylbutyl)-L- $\alpha$ -aspartyl]-L-phenylalanine 1-methyl ester, the concentration depending of course on the desired end use.

The Examples which follow are intended as an illustration of certain preferred embodiments of the invention, and no limitation of the invention is implied.

25

#### Example 1

Hydrochloride Salt of N-[N-(3,3-dimethylbutyl)-L- $\alpha$ -aspartyl]-L-phenylalanine 1-methyl ester

N-[N-(3,3-dimethylbutyl)-L- $\alpha$ -aspartyl]-L-phenylalanine 1-methyl ester (25.0 g) was slurried in 150 ml of water. To the stirred slurry was slowly added concentrated hydrochloric acid (5.5 ml, 0.0661 mol). After approximately 10 minutes all the solid had dissolved. The clear solution was then freeze dried. The resulting product exhibited a greatly increased aqueous solubility compared to the starting material. Yield: 27.34 g white solid (99.7%). The hydrochloride

- 8 -

salt (1 g) dissolved in water (100 ml) in 70 seconds. <sup>1</sup>H NMR (CD<sub>3</sub>CN) δ 0.89 (s, 9H), 1.64 (m, 2H), 3.01 (m, 6H), 3.68 (s, 3H), 4.16 (t, 1H), 4.67 (m, 1H), 7.26 (m, 5H), 8.27 (d, 1H). Anal. Calcd for C<sub>20</sub>H<sub>31</sub>ClN<sub>2</sub>O<sub>5</sub>•H<sub>2</sub>O: C, 55.48; H, 7.69; N, 6.47; Cl, 8.19. Found: C, 55.95; H, 7.59; N, 6.55; Cl, 8.35.

## Example 2

10           Phosphate Salt of N-[N-(3,3-dimethylbutyl)-  
L-α-aspartyl]-L-phenylalanine 1-methyl ester

N-[N-(3,3-dimethylbutyl)-L-α-aspartyl]-L-phenylalanine 1-methyl ester (10.0 g) was slurried in 100 ml of water. An equimolar amount of 85% phosphoric acid solution was added. After all the solid dissolved, the solution was freeze dried. The resulting product exhibited a greatly increased aqueous solubility compared to the starting material. Yield 12.33 g white solid (98.0%). The phosphate salt (0.1 g) dissolved in water (100 ml) in less than 30 seconds (visual observation) and about 90 seconds by spectrophotometric analysis. <sup>1</sup>H NMR (CD<sub>3</sub>CN) δ 0.89 (s, 9H), 1.55 (m, 2H), 2.84 (m, 5H), 3.19 (m, 1H), 3.67 (s, 3H), 4.06 (t, 1H), 4.70 (m, 1H), 7.27 (m, 5H), 8.03 (d, 1H). Anal. Calcd for C<sub>20</sub>H<sub>33</sub>N<sub>2</sub>O<sub>9</sub>P•H<sub>2</sub>O: C, 48.58; H, 7.15; N, 5.67; P, 6.26. Found: C, 48.15; H, 7.08; N, 5.60; P, 6.57.

## Example 3

30           Sulfate Salt of N-[N-(3,3-dimethylbutyl)-  
L-α-aspartyl]-L-phenylalanine 1-methyl ester

N-[N-(3,3-dimethylbutyl)-L-α-aspartyl]-L-phenylalanine 1-methyl ester (25.0 g) was slurried in 100 ml of water and concentrated sulfuric acid (3.67 ml) was added slowly. After several minutes of stirring some material remained undissolved. Therefore, 75 ml of acetone was added. The resulting clear solution was stirred for 30 minutes, then the acetone was removed

under reduced pressure on a rotary evaporator. The contents of the flask remained a clear solution. Freeze drying of this solution afforded a white solid which exhibits a greater aqueous solubility compared to  
5 the starting material. Yield: 30.04 g (95%). The sulfate salt (0.1 g) dissolved in water (100 ml) in less than 20 seconds (visual observation). <sup>1</sup>H NMR (CD<sub>3</sub>CN) δ 0.89 (s, 9H), 1.57 (m, 2H), 2.91 (m, 5H), 3.20 (m, 1H), 3.68 (s, 3H), 4.15 (t, 1H), 4.72 (m, 1H), 7.27  
10 (m, 5H), 7.68 (d, 1H). Anal. Calcd for C<sub>20</sub>H<sub>32</sub>N<sub>2</sub>O<sub>9</sub>S•H<sub>2</sub>O: C, 48.56; H, 6.94; N, 5.66; S, 6.48. Found C, 48.93; H, 7.02; N, 5.78; S, 6.50.

#### Example 4

15 Citrate Salt of N-[N-(3,3-dimethylbutyl)-L-α-aspartyl]-L-phenylalanine 1-methyl ester

N-[N-(3,3-dimethylbutyl)-L-α-aspartyl]-L-phenylalanine 1-methyl ester (5.00g) was dissolved in 75 ml of  
20 acetone. Citric acid (2.53 g) was dissolved in 50 ml of water, and the acetone solution was added. This formed a clear, colorless solution that was stirred for 1 hour. The acetone was evaporated under reduced pressure. The resulting slurry was freeze dried to  
25 obtain 7.5 g of a white solid. The citrate salt (0.1 g) dissolved in water (100 ml) in 90 seconds (visual observation). δ (CD<sub>3</sub>CN) 0.89 (s, 9H), 1.53 (m, 2H), 2.83 (m, 10H), 3.20 (m, 1H), 3.67 (s, 3H), 4.04 (t, 1H), 4.21 (m, 1H), 7.27 (m, 5H), 7.79 (d, 1H). Anal.  
30 Calcd for C<sub>26</sub>H<sub>38</sub>N<sub>2</sub>O<sub>12</sub>•H<sub>2</sub>O: C, 53.05; H, 6.86; N, 4.76. Found: C, 53.10; H, 6.88; N, 4.41.

#### Comparative Example 1

35 Dissolution of N-[N-(3,3-dimethylbutyl)-L-α-aspartyl]-L-phenylalanine 1-methyl ester in Water

N-[N-(3,3-dimethylbutyl)-L-α-aspartyl]-L-phenylalanine 1-methyl ester (0.05 - 0.1 g) was dissolved in water


- 10 -

(100 mL). The compound completely dissolved in 5-7 minutes (visual observation). The dissolution of 1.0 g of N-[N-(3,3-dimethylbutyl)-L- $\alpha$ -aspartyl]-L-phenylalanine 1-methyl ester in 100 mL of water  
5 required approximately 45 minutes.

Other variations and modifications of this invention will be obvious to those skilled in this art. This  
10 invention is not to be limited except as set forth in the following claims.

WHAT IS CLAIMED IS:

1. A salt of N-[N-(3,3-dimethylbutyl)-L- $\alpha$ -aspartyl]-L-phenylalanine 1-methyl ester represented by the formula



wherein  $X^-$  is selected from the group consisting of  $Cl^-$ ,  $HSO_4^-$ ,  $H_2PO_3^-$ , citrate $^-$ ,  $HCO_3^-$ , fumarate, malate, maleate, tartarate, acetate, benzoate or gluconate.


2. A salt according to claim 1, wherein  $X^-$  is  $Cl^-$ .
3. A salt according to claim 1, wherein  $X^-$  is  $HSO_4^-$ .
4. A salt according to claim 1, wherein  $X^-$  is  $H_2PO_3^-$ .
5. A salt according to claim 1, wherein  $X^-$  is citrate.
6. A salt according to claim 1, wherein  $X^-$  is  $HCO_3^-$ .
7. A salt according to claim 1, wherein  $X^-$  is fumarate.
8. A salt according to claim 1, wherein  $X^-$  is malate.
9. A salt according to claim 1, wherein  $X^-$  is maleate.
10. A salt according to claim 1, wherein  $X^-$  is tartarate.
11. A salt according to claim 1, wherein  $X^-$  is acetate.

- 12 -

12. A salt according to claim 1, wherein  $X^-$  is benzoate.

13. A salt according to claim 1, wherein  $X^-$  is gluconate.

14. A liquid low-calorie sweetener composition comprising a salt of a dipeptide-sweetener represented by the formula



wherein  $X^-$  is selected from the group consisting of  $Cl^-$ ,  $HSO_4^-$ ,  $H_2PO_3^-$ , citrate<sup>-</sup>,  $HCO_3^-$ , fumarate, malate, maleate, tartarate, acetate, benzoate or gluconate, dissolved in a consumable solvent or solvents in a concentration up to about 40% by weight of the composition to provide a high concentration liquid low-calorie sweetener.

15. A liquid low-calorie sweetener according to claim 14, wherein the solvent is ethanol, water, propylene glycol or a mixture thereof.

A. CLASSIFICATION OF SUBJECT MATTER  
 IPC 6 C07K5/06 A23L1/236

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07K A23L

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	FR 2 697 844 A (NOFRE & TINTI) 13 May 1994 see the whole document & US 5 480 668 A (NOFRE & TINTI) cited in the application	1-15
Y	FR 2 182 921 A (GENERAL FOODS CORPORATION) 14 December 1973 see the whole document & US 4 029 701 A (GENERAL FOODS CORPORATION) cited in the application	1-15

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

\* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

2 February 1999

Date of mailing of the international search report


10/02/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
 NL - 2280 HV Rijswijk  
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
 Fax: (+31-70) 340-3016

Authorized officer

Masturzo, P





Patent document cited in search report	Publication date	Patent family member(s)	Publication date
FR 2697844 A	13-05-1994	AT 138935 T	15-06-1996
		AU 664663 B	23-11-1995
		AU 5468194 A	08-06-1994
		BG 61609 B	30-01-1998
		BG 99299 A	29-09-1995
		CA 2139233 A	26-05-1994
		CN 1090571 A,B	10-08-1994
		CZ 9403319 A	18-10-1995
		DE 69303032 D	11-07-1996
		DE 69303032 T	02-10-1996
		DK 669935 T	01-07-1996
		EP 0669935 A	06-09-1995
		ES 2091114 T	16-10-1996
		FI 945451 A	22-12-1994
		WO 9411391 A	26-05-1994
		GR 3020164 T	30-09-1996
		HU 72192 A	28-03-1996
		JP 2818032 B	30-10-1998
		JP 8503206 T	09-04-1996
		LT 1457 A,B	15-06-1994
		MD 960256 A	31-03-1998
		NO 945090 A	30-12-1994
		NZ 257870 A	26-03-1996
		PL 306841 A	18-04-1995
		SK 158694 A	10-05-1995
		US 5480668 A	02-01-1996
		ZA 9308430 A	13-06-1994
FR 2182921 A	14-12-1973	US 4029701 A	14-06-1977
		US 4031258 A	21-06-1977
		CA 1028195 A	21-03-1978
		DE 2315646 A	11-10-1973
		GB 1431057 A	07-04-1976
		JP 1061651 C	31-08-1981
		JP 49013371 A	05-02-1974
		JP 56003015 B	22-01-1981
		NL 7304314 A	02-10-1973
		CA 1028319 A	21-03-1978

